

MORSE POSITION OF KNOTS AND CLOSED INCOMPRESSIBLE SURFACES

MAKOTO OZAWA

ABSTRACT. In this paper, we study on knots and closed incompressible surfaces in the 3-sphere via Morse functions. We show that both of knots and closed incompressible surfaces can be isotoped into a "related Morse position" simultaneously. As an application, we have following results.

- Smallness of Montesinos tangles with length two and Kinoshita's theta curve
- Classification of closed incompressible and meridionally incompressible surfaces in 2-bridge theta-curve and handcuff graph complements and the complements of links which admit Hopf tangle decompositions

1. INTRODUCTION

Morse theory had a tremendous impact on low-dimensional topology. Bridge positions of knots that developed by Schubert ([7]) and Heegaard splittings of 3-manifolds that developed by Heegaard ([3]) are independent of Morse theory, but those can be explained integrally by Morse theory.

A bridge position of a knot is defined by using a Morse function from the 3-sphere to the 1-dimensional Euclidean space with two critical points so that any maximal critical point of the knot is situated above any minimal critical point. This general idea was expanded to a notion of thin position by Gabai, and he settled the Property R conjecture ([1]). Thin position also brought a celebrated solution of the knot complement conjecture by Gordon and Luecke ([2]).

In this paper, we show that we can isotope a closed incompressible surface to a related Morse position in the complement of a knot placed in a bridge or thin position. Then as an applications, we show smallness of Montesinos tangles with length two and Kinoshita's theta curve, and classification of closed incompressible and meridionally incompressible surfaces in 2-bridge theta-curve and handcuff graph complements and the complements of links which admit Hopf tangle decompositions.

2. RESULT

Let K be a knot in the 3-sphere S^3 and $h : S^3 \rightarrow \mathbb{R}$ be the composition of the inclusion $S^3 \subset \mathbb{R}^4$ and the height function $\mathbb{R}^4 \rightarrow \mathbb{R}$ defined by $(x, y, z, w) \mapsto w$. We say that K is in a *Morse position* with respect to h if the restriction of h to K is a Morse function. Let $a_0, \dots, a_n$ be the critical points of K which are labeled so that the corresponding critical values $t_i = h(a_i)$ satisfy $t_{i-1} < t_i$ for all i . For

1991 *Mathematics Subject Classification*. Primary 57M25, 57M50; Secondary 57Q35, 57Q37.

Key words and phrases. Knot, Morse position, Bridge position, Thin position, Incompressible surface.

a regular value $s_i \in (t_{i-1}, t_i)$, $P_i = h^{-1}(s_i)$ is called a *level sphere* between a_{i-1} and a_i . The *width* of K is $w(K) = \sum_{i=1}^n |P_i \cap K|$. A knot K is in *thin position* if $w(K)$ is minimal up to isotopy of K . A level sphere P_i between a_{i-1} and a_i is said to be *thin* if a_{i-1} is a local maximum and a_i a local minimum. Similarly, a level sphere P_i between a_{i-1} and a_i is said to be *thick* if a_{i-1} is a local minimum and a_i a local maximum. A knot K is in *bridge position* if there is only one thick level sphere $P_{\frac{n+1}{2}}$. Then, the *bridge number* of K is $b(K) = |P_{\frac{n+1}{2}} \cap K|/2$. A knot K is in *minimal bridge position* if $b(K)$ is minimal up to isotopy of K . For each thick level sphere P_i between a_{i-1} and a_i , we define the *thick region* of K as $h^{-1}([t_{i-1} + \epsilon, t_i - \epsilon])$ for a sufficiently small positive real number ϵ . Then, no thick region contains a maximum or minimum of K , and the rest of all thick regions in S^3 contain all maxima and minima of K , each of which is called a *thin region* of K . Any thick region contains only one thick level sphere, and any thin region, except for the top 3-ball and the bottom 3-ball, contains only one thin level sphere. Let M_{thin} be the disjoint union of all thin regions of K , and M_{thick} the disjoint union of all thick regions of K .

Let F be a closed surface in S^3 disjoint from K or intersecting K transversely. We say that F is in a *Morse position* with respect to h if the restriction of h to F is a Morse function. The critical points of K and F are assumed to be mutually disjoint. Moreover, we say that F is in a *Morse position related to K* if the following conditions are satisfied.

- (1) F is in a Morse position with respect to h ,
- (2) F and K are in general position and $F \cap K$ is contained in M_{thick} ,
- (3) all maxima and minima of F are contained in M_{thin} , and all saddles of F are contained in M_{thick} .

These three conditions are easily satisfied for any closed surface F . We prepare definitions to describe the next condition (4).

Let M be a 3-manifold, T a 1-manifold properly embedded in M , and F a surface properly embedded in M such that F is disjoint from T or intersects T transversely in the interior of F . An *isotopy* ϕ_t of F in (M, T) should be assumed that $\phi_t(F)$ and T are in general position for all $t \in [0, 1]$. We recall the definitions about incompressible surfaces and meridionally incompressible surfaces.

First, suppose that there exists a disk D in $M - T$ such that $D \cap F = \partial D$ and ∂D is essential in $F - T$. Then by cutting F along ∂D and pasting two parallel copies of D , we have a new surface F' . We say that F' is obtained from F by *compressing F along D* , and D is called a *compressing disk* for F . A surface F is *incompressible* in (M, T) if for any disk D in $M - T$ with $D \cap F = \partial D$, there exists a disk D' in $F - T$ such that $\partial D' = \partial D$. In the case that F is a disk disjoint from T , F is said to be *incompressible* in (M, T) if for any disk D in $\partial M - \partial T$ with $\partial D = \partial F$, $F \cup D$ does not bound a 3-ball in $M - T$. In the case that F is a 2-sphere disjoint from T , F is said to be *incompressible* in (M, T) if F does not bound a 3-ball in $M - T$.

Next, suppose that there exists a disk D in M such that $D \cap F = \partial D$ and D intersects T in one point of the interior of D . Then by cutting F along ∂D and pasting two parallel copies of D , we have a new surface F' . If F is incompressible in (M, T) , then F' is also incompressible in (M, T) . We say that F is *meridionally incompressible* in (M, T) if for any disk D in M such that $D \cap F = \partial D$ and D intersects T in one point of the interior of D , there exists a disk D' in F such that $\partial D' = \partial D$ and $D \cup D'$ bounds a trivial (3-ball, arc)-pair in (M, T) . In the case

that F is a disk intersecting T in one point in the interior of F , we say that F is *meridionally incompressible* in (M, T) if for any disk D in ∂M with $\partial D = \partial F$, $F \cup D$ does not bound a trivial (3-ball, arc)-pair in (M, T) . In the case that F is a 2-sphere intersecting T in two points, we say that F is *meridionally incompressible* in (M, T) if F does not bound a trivial (3-ball, arc)-pair in (M, T) . We note that this definition is slightly different from a usual one as a matter of convenience.

Suppose that $K \subset S^3$ is a knot in a Morse position with respect to h and $F \subset S^3$ is a closed surface in a Morse position related to K . Furthermore, we say that F is in an *essential Morse position related to K* if the following condition is satisfied.

- (4) each component of $F \cap M_{thin}$ and $F \cap M_{thick}$ is incompressible and meridionally incompressible in $(M_{thin}, K \cap M_{thin})$ and $(M_{thick}, K \cap M_{thick})$ respectively.

The following main theorem has a philosophy that makes thin regions M_{thin} simple, but thick regions M_{thick} essential.

Theorem 2.1. *Let $K \subset S^3$ be a knot in a Morse position with respect to h and $F \subset S^3$ be a closed surface disjoint from K or intersecting K transversely such that F is incompressible and meridionally incompressible in (S^3, K) . Then F can be isotoped so that;*

- (1) F is a thin level sphere, or
- (2) F is in an essential Morse position related to K .

Example 2.2. Let K be a composite knot of two trefoils, and F be the decomposing sphere for K . If we put K in a bridge position, then by Theorem 2.1, F can be isotoped so that F is in an essential Morse position related to K . See Figure 1.




FIGURE 1. Simple example of type 2 in Theorem 2.1

If we put K in a thin position, then by Theorem 2.1, F can be isotoped so that F is in a thin level sphere. See Figure 2.





FIGURE 2. Simple example of type 1 in Theorem 2.1

A 3-manifold M is said to be *small* if any closed incompressible surface properly embedded in M is ∂ -parallel. Analogously, a knot $K \subset S^3$ is *small* if the exterior $E(K) = S^3 - \text{int}N(K)$ is small, a tangle (B, T) is *small* if $B - \text{int}N(T)$ is small, and a graph $G \subset S^3$ is *small* if $S^3 - \text{int}N(G)$ is small.

A *Montesinos tangle* $T(r_1, \dots, r_n)$ is defined as a series of n rational tangles of slope r_i ($i = 1, \dots, n$). See Figure 3. We note that if $r_i = \infty$ for some i , then $T(r_1, \dots, r_n)$ is splittable by a properly embedded disk, and that if r_i is an integer, then the length n of $T(r_1, \dots, r_n)$ can be taken short. For this reason, it is reasonable to assume that $r_i \neq \infty$ and r_i is not an integer for all i .

FIGURE 3. Montesinos tangle $T(r_1, \dots, r_n)$

Theorem 2.3. *A Montesinos tangle $T(r_1, r_2)$ is small if $r_i \neq \infty$ for $i = 1, 2$.*

Remark 2.4. The referee suggested the possibility to show that a tangle is small by showing the double branched cover is small. Indeed, it seems true and Theorem 2.3 follows the smallness of the double branched cover over a Montesinos tangle $T(r_1, r_2)$. Theorem 2.3 gives an another direct proof without branched covering spaces or orbifolds ([6, Theorem 1]).

Example 2.5. As a corollary of Theorem 2.3, we can show that Kinoshita's theta curve is small. Kinoshita's theta curve is known as a most popular theta curve, which is normally illustrated as Figure 4. It is known to be minimally knotted and hyperbolic. Therefore, the exterior is ∂ -irreducible.




FIGURE 4. standard form of Kinoshita's theta curve

We shorten an edge and obtain an another form of Kinoshita's theta curve as in Figure 5. Then, the 2-string tangle on the left-hand side is a Montesinos tangle with length two and the right-hand side is a trivial H -shaped graph tangle. Hence, the exterior of Kinoshita's theta curve is homeomorphic to the exterior of the Montesinos tangle. This fact was also observed by Wu ([8, Example 3]). Therefore, Kinoshita's theta curve is small by Theorem 2.3.




FIGURE 5. another form of Kinoshita's theta curve

Let G be a theta curve or handcuff graph embedded in S^3 . We say that G is in a *Morse position* with respect to h if the restriction of h to each edge of G is a Morse function. In this paper, we assume that near each vertex v of G , three edges either go up or go down from v . We say that a theta curve or handcuff graph G in a Morse position is *2-bridge* if there exists a level 2-sphere P which intersects G in 5 points other than vertices and critical points, and edges of G have exactly two critical points.

Let $F \subset S^3$ be a closed surface disjoint from G or intersecting edges of G transversely. Concerning the definition for F to be meridionally incompressible, we say that F is *meridionally incompressible* in (S^3, G) if F is not parallel to $\partial N(v)$ for a vertex v of G , together with the definition for knot case.

The next theorem asserts there is no incompressible and meridionally incompressible surface of positive genus in the complement of 2-bridge theta curves.

Theorem 2.6. *Let G be a 2-bridge theta curve or handcuff graph in S^3 , and F be an incompressible and meridionally incompressible surface. Then F is isotopic to a sphere with one maximum and one minimum illustrated in Figure 6.*




FIGURE 6. incompressible and meridionally incompressible surface in 2-bridge theta curve or handcuff graph complements

As a corollary of Theorem 2.6, we obtain the Motohashi's result ([5]) about composite 2-bridge theta curves. A theta curve G is *composite* if there exists a 2-sphere which intersects each edge of G in one point and decompose G into two non-trivial theta curve. By Theorem 2.6, the decomposing sphere is isotopic to a

sphere with one maximum and minimum as Figure 7. Hence, G is decomposed into two rational theta curves, that is, a theta curve obtained by pasting a rational tangle and a trivial H -shaped graph tangle.




FIGURE 7. composite 2-bridge theta curve and the decomposing sphere

Next, we consider closed incompressible surfaces in the complements of links that admit Hopf tangle decompositions. The *Hopf tangle* (B, T) is a 2-string trivial tangle $(B, t_1 \cup t_2)$ with a trivial loop C as illustrated as Figure 8. The *equator* of the Hopf tangle (B, T) is defined as a loop e in ∂B which cobounds an annulus in $B - (t_1 \cup t_2)$ with the trivial loop C , and the *meridian* is defined as a loop m in ∂B which bounds a disk D in $B - (t_1 \cup t_2)$ separating two strings of T .




FIGURE 8. Hopf tangle (B, T)

Let $\mathcal{L}(x_1, x_2)$ be the set of links which are obtained by gluing two Hopf tangles (B_1, T_1) and (B_2, T_2) so that x_1 coincides with x_2 , where x_i represents either the equator e_i of (B_i, T_i) or the meridian m_i of (B_i, T_i) for $i = 1, 2$.

We note that if a link L in S^3 admits a Hopf tangle decomposition $(S^3, L) = (B_1, T_1) \cup (B_2, T_2)$, then the decomposing sphere $S = \partial B_1 = \partial B_2$ is incompressible and meridionally incompressible in (S^3, L) .

Remark 2.7. We remark that $\mathcal{L}(m_1, m_2) \cap (\mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2)) = \emptyset$ and $(\mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2)) \cap \mathcal{L}(e_1, e_2) = \emptyset$, and that $\mathcal{L}(m_1, m_2) \cap \mathcal{L}(e_1, e_2)$ consists of one link which is obtained by the orientation reversing identity map between two Hopf tangles, namely, the Borromean rings.

Theorem 2.8. *Let L be a link in S^3 which admits a Hopf tangle decomposition $(S^3, L) = (B_1, T_1) \cup (B_2, T_2)$. Then,*

- (1) *There exists an incompressible and meridionally incompressible closed surface in (S^3, L) different from the decomposing sphere if and only if $L \in \mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2) \cup \mathcal{L}(e_1, e_2)$.*
- (2) *L admits a unique Hopf tangle decomposition if and only if $L \notin \mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2)$. If $L \in \mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2)$ except the Borromean rings, then L admits exactly two Hopf tangle decompositions. The Borromean rings admits exactly three Hopf tangle decompositions.*
- (3) *L is small if and only if $L \notin \mathcal{L}(e_1, e_2)$. If $L \in \mathcal{L}(e_1, e_2)$, then there exists an essential torus in $S^3 - \text{int}N(L)$.*

See Figure 9 for the types of link classes and associated closed incompressible and meridionally incompressible surfaces, where dotted lines represent monotone strings (which may have some twists).




FIGURE 9. three types of link classes and closed incompressible and meridionally incompressible surfaces

Example 2.9. The Borromean rings is the only element of $\mathcal{L}(m_1, e_2) \cap \mathcal{L}(e_1, m_2)$ since it can be deformed as Figure 10. By Remark 2.7, it does not belong to $\mathcal{L}(m_1, m_2)$ and $\mathcal{L}(e_1, e_2)$. Theorem 2.8 shows that the Borromean rings has exactly

three Hopf tangle decomposing spheres F_1 , F_2 and F_3 , and that it is small. Matsuda ([4]) also proved that the Borromean rings is small by using clasp disks bounded by the Borromean rings.




FIGURE 10. the Borromean rings and three Hopf tangle decomposing spheres

3. PRELIMINARY

Let S be a 2-sphere and T a 1-manifold properly embedded in $S \times I$. In this section, we consider incompressible surfaces embedded in $(S \times I, T)$, with the projection $p : S \times I \rightarrow S \times \{0\}$ and the height function $h : S \times I \rightarrow I$. Let F be a surface properly embedded in $S \times I$ such that F is disjoint from T or intersects T transversely in the interior of F . An isotopy ϕ_t of $S \times I$ is called a *horizontal isotopy* if $h \circ \phi_t = h$ for all $t \in [0, 1]$. A surface F is said to be *horizontally ∂ -parallel* in $S \times I$ if F is ∂ -parallel in $S \times I$ and there exists a horizontal isotopy ϕ_t of $S \times I$ such that $p \circ \phi_1$ is a homeomorphism on F . We say that F is *T -compressible* in $(S \times I, T)$ if there exists a disk D in $S \times I$ such that $D \cap T = \partial D \cap T = \alpha$ is a subarc of T and $D \cap F = \partial D - \text{int} \alpha$. A surface F is *T -incompressible* in $(S \times I, T)$ if it is not T -compressible.

An n -dimensional submanifold V in $S \times I$ is *vertical* if $p(V)$ is an $(n - 1)$ -dimensional manifold. A submanifold H in $S \times I$ is *horizontal* if H is contained in a level 2-sphere $S \times \{t\}$ for some $t \in I$.

Lemma 3.1. *Let S be a 2-sphere, X a union of points in S , and $F \subset (S \times I, X \times I)$ an $(X \times I)$ -incompressible surface. If F is horizontally ∂ -parallel in $S \times I$, then F is ∂ -parallel in $(S \times I, X \times I)$.*

Proof. After a suitable horizontal isotopy of $S \times I$, we may assume that p is a homeomorphism on F . Moreover, by a suitable horizontal isotopy of $S \times I$ fixing F , we may assume that p is a homeomorphism on $(X \times I) \cap V$ since $X \times I$ is

monotone in $S \times I$, where V is a 3-manifold bounded by F and $p(F)$ in $S \times I$. Then $p((X \times I) \cap V)$ consists of disjoint arcs embedded in $p(F)$. See Figure 11.




FIGURE 11. F is α -compressible

Suppose that there exists an arc α of $(X \times I) \cap V$ such that $\partial\alpha$ is contained in F . We note that the vertical disk $p^{-1}(p(\alpha)) \cap V$ is disjoint from $((X \times I) \cap V) - \alpha$. Then, there exists a disk D in $p^{-1}(p(\alpha)) \cap V$ such that $D \cap \alpha = \partial D \cap \alpha = \alpha$ and $\partial D - \text{int}\alpha \subset F$. This shows that F is an $(X \times I)$ -compressible in $(S \times I, X \times I)$, a contradiction. Hence, any arc of $(X \times I) \cap V$ connects F and $p(F)$ monotonously, and F is ∂ -parallel in $(S \times I, X \times I)$. $\square$

Lemma 3.2. *Let S be a 2-sphere, X a union of points in S , and $F \subset (S \times I, X \times I)$ an incompressible and meridionally incompressible surface in a Morse position with respect to h . Suppose that the number of critical points of F is minimal up to isotopy of F in $(S \times I, X \times I)$. If F has a maximal critical point, then F is ∂ -parallel in $(S \times I, X \times I)$.*

Proof. If F is horizontally ∂ -parallel in $S \times I$, then by Lemma 3.1, F is ∂ -parallel in $(S \times I, X \times I)$. So, hereafter we assume that F is not horizontally ∂ -parallel in $S \times I$.

Let $a_0, \dots, a_n$ be the critical points of F which are labeled so that the corresponding critical values $t_i = h(a_i)$ satisfy $t_{i-1} < t_i$ for all i , and let a_m be the lowest

maximal critical point of F . If $m = 0$, then F is a disk with only one maximal critical point a_0 , and hence horizontally ∂ -parallel in $S \times I$. This contradicts the assumption. Hence, we have $m \geq 1$.

Let F_{max} be the maximal horizontally ∂ -parallel subsurface of F containing a_m , that is, a component of $F \cap h^{-1}([t_{m-j} + \epsilon, t_m + \epsilon])$ which is horizontally ∂ -parallel in $h^{-1}([t_{m-j} + \epsilon, t_m + \epsilon])$, for an integer j ($\leq m$) as large as possible and a fixed sufficiently small positive real number ϵ . See Figure 12.




FIGURE 12. maximal horizontally ∂ -parallel subsurface F_{max}

Since F_{max} is horizontally ∂ -parallel in $S \times [t_{m-j} + \epsilon, t_m + \epsilon]$ and $(X \times [t_{m-j} + \epsilon, t_m + \epsilon])$ -incompressible in $(S \times [t_{m-j} + \epsilon, t_m + \epsilon], X \times [t_{m-j} + \epsilon, t_m + \epsilon])$, by Lemma 3.1, it is a ∂ -parallel planar surface in $(S \times [t_{m-j} + \epsilon, t_m + \epsilon], X \times [t_{m-j} + \epsilon, t_m + \epsilon])$. Therefore, by a suitable horizontal isotopy of $S \times I$, we may assume that p is a homeomorphism on F_{max} . Then, F_{max} is isotopic to $F'_{max} = p^{-1}(p(F_{max})) \cap (S \times \{t_{m-j} + \epsilon\})$ in $(S \times [t_{m-j} + \epsilon, t_m + \epsilon], X \times [t_{m-j} + \epsilon, t_m + \epsilon])$, and F_{max} and F'_{max} bound a submanifold V in $S \times [t_{m-j} + \epsilon, t_m + \epsilon]$ such that $p(F_{max}) = p(F'_{max}) = p(V)$. We note that $\text{int} V \cap F = \emptyset$ since a_m is the lowest maximal critical point of F . Moreover, by the minimality of the number of critical points of F , F_{max} has only one maximal point a_m and $(|\partial F_{max}| - 1)$ -saddle points.

Hereafter, we consider the critical point a_{m-j} . If a_{m-j} is a saddle point, then the saddle transformation at a_{m-j} can be regarded as attaching a horizontal rectangular band B to $F \cap (S \times [t_{m-j} + \epsilon, t_m + \epsilon])$ in a usual way. Otherwise, since a_m is the lowest maximal point, a_{m-j} is a minimal point and can be regarded as attaching a horizontal disk E to $F \cap (S \times [t_{m-j} + \epsilon, t_m + \epsilon])$. Then, there are the following possibility for the critical point a_{m-j} .

Case 1: The band B is outside of F'_{max} and connects a same boundary component of F_{max} .

Case 2: The band B is outside of F'_{max} and connects a boundary component of F_{max} and a component of $F \cap (S \times [t_{m-j} + \epsilon, t_m + \epsilon])$ other than F_{max} .

Case 3: The band B is inside of F'_{max} and connects a same boundary component of F_{max} .

Case 4: The band B is inside of F'_{max} and connects two different boundary components of F_{max} .

Case 5: The disk E caps F_{max} along a boundary component of F_{max} .

In Case 1, p is a homeomorphism on $F_{max} \cup B$. Hence, a component of $F \cap h^{-1}([t_{m-j} - \epsilon, t_m + \epsilon])$ containing a_m is horizontally ∂ -parallel in $h^{-1}([t_{m-j} - \epsilon, t_m + \epsilon])$. This contradicts the maximality of F_{max} .

In Case 2, by an isotopy of F_{max} to F'_{max} and pushing out it through F'_{max} , the maximal critical point a_m and the saddle point a_{m-j} are mutually canceled without increasing the number of critical points of F . This contradicts the minimality of the number of critical points of F .

In Case 3, there exists a vertical compressing disk D for $F_{max} \cup B$ in $(S \times [t_{m-j} - \epsilon, t_m + \epsilon], X \times [t_{m-j} - \epsilon, t_m + \epsilon])$ such that $D \cap (X \times I) = \emptyset$, $D \cap B = \partial D \cap B = \alpha$ is an arc through B , and $p(D) = p(\alpha)$. By the incompressibility of F in $(S \times I, X \times I)$, there exists a disk D' in F such that $\partial D' = \partial D$. Then, by an isotopy of D' to D and a suitable isotopy of F , the saddle point a_{m-j} is eliminated without increasing the number of critical points of F . This contradicts the minimality of the number of critical points of F .

In Case 4, there exists a vertical compressing disk D for $F_{max} \cup B$, as same as Case 3. Actually, this disk D gives a compressing disk for F in $(S \times I, X \times I)$ since ∂D is essential in F . This contradicts the incompressibility of F in $(S \times I, X \times I)$.

In Case 5, first we consider the case that F_{max} is a disk. If $p(F_{max})$ contains $p(E)$, then $F = F_{max} \cup E$ is a 2-sphere which bounds a 3-ball V' in $S \times I$. We note that $V' \cap (X \times I) = \emptyset$ since $E \cap (X \times I) = \emptyset$ and F_{max} is isotopic to E in $(V', (X \times I) \cap V')$. Hence F bounds a 3-ball V' in $S \times I - X \times I$, and this contradicts the incompressibility of F in $(S \times I, X \times I)$. Otherwise, $F_{max} \cup E$ is isotopic to a level 2-sphere $h^{-1}(t_{m-j})$ in $(S \times I, X \times I)$, but this contradicts the assumption that F is not horizontally ∂ -parallel in $S \times I$.

Next, we consider the case that F_{max} is not a disk. In this case, $p(F_{max})$ does not contain $p(E)$. Hence, $F_{max} \cup E$ is isotopic to a planar surface in $h^{-1}(t_{m-j})$, and there exists an isotopy of $F_{max} \cup E$ so that it has only one maximal point a_m and $(|\partial F_{max}| - 2)$ -saddle points since a saddle point of F_{max} and a minimal point a_{m-j} are mutually canceled. This contradicts the minimality of the number of critical points of F .

Hence, any case does not occur, the lemma is proved. $\square$

The next lemma is a special case of Lemma 3.2. In the case that the number of strings is less than or equal to 5, incompressible and meridionally incompressible surfaces are very restricted.

Lemma 3.3. *Let S be a 2-sphere, X a union of points in S less than or equal to 5, and $F \subset (S \times I, X \times I)$ an incompressible and meridionally incompressible surface in a Morse position with respect to h . Suppose that the number of critical points of F is minimal up to isotopy of F in $(S \times I, X \times I)$. Then, one of the following holds.*

- (1) *F has no critical point and any loop of $F \cap (S \times \{t\})$ does not bound a disk D in $S \times \{t\}$ such that $|D \cap (X \times I)| \leq 1$.*

(2) F is ∂ -parallel in $(S \times I, X \times I)$.

Proof. If F has a maximal or minimal critical point, then by Lemma 3.2, F is ∂ -parallel in $(S \times I, X \times I)$ and we have a conclusion 2 of Lemma 3.3.

Hereafter, we assume that F has neither a maximal nor minimal critical point, and suppose that F has a saddle point a_s . Let F_s be a component of $F \cap h^{-1}([h(a_s) - \epsilon, h(a_s) + \epsilon])$ containing a_s . Then, at least one of boundary components of F_s bounds a disk in $S \times \{h(a_s) \pm \epsilon\}$ which is disjoint from $X \times I$ or intersects $X \times I$ in one point. Therefore, there exists a horizontal compressing disk or horizontal meridionally compressing disk D for F in $(S \times [h(a_s) - \epsilon, h(a_s) + \epsilon], X \times [h(a_s) - \epsilon, h(a_s) + \epsilon])$. Since F is incompressible and meridionally incompressible in $(S \times I, X \times I)$, there exists a disk D' in F such that $\partial D' = \partial D$ and $D \cup D'$ bounds a 3-ball in $S \times I - X \times I$ or a trivial (3-ball, arc)-pair in $(S \times I, X \times I)$. Moreover, since D is horizontal, D' contains at least one maximal or minimal critical point. This contradicts the assumption and hence F has no critical point. Similarly, if there exists a loop of $F \cap (S \times \{t\})$ which bounds a disk D in $S \times \{t\}$ such that $|D \cap (X \times I)| \leq 1$, then it contradicts the assumption. Therefore, we have a conclusion 1 of Lemma 3.3. $\square$

Lemma 3.4. *Let (B, T) be a rational tangle, F a surface properly embedded in B which is disjoint from T or intersects T transversely. If F is incompressible and meridionally incompressible in (B, T) , then one of the following holds.*

- (1) F is ∂ -parallel in (B, T) .
- (2) F is a disk separating two strings of T .

Proof. Let B be the north hemisphere $\{(x, y, z, w) \in \mathbb{R}^4 | x^2 + y^2 + z^2 + w^2 = 1, w \geq 0\}$, and $h : B \rightarrow \mathbb{R}$ be the height function defined by $(x, y, z, w) \mapsto w$. Since (B, T) is a rational tangle, T can be isotoped so that the restriction of h to T is a Morse function with two maximal critical points. Let S a level 2-sphere below two maximal points of T , and put $M_{thin} = B \cap h^{-1}([h(S), 1])$ and $M_{thick} = B \cap h^{-1}([0, h(S)])$. See Figure 13.




FIGURE 13. M_{thin} - M_{thick} decomposition of (B, T)

We isotope F so that F is a Morse position with respect to h . In the same way as Claim 4.1, we may assume that $F \cap M_{thin}$ consists of incompressible disks which separate two strings of $T \cap M_{thin}$. Hereafter, we suppose that $|F \cap M_{thin}|$ and the critical points of F are minimal up to isotopy of F in (B, T) . Then each component of $F \cap M_{thin}$ is an incompressible disk with only one maximal critical point, and by the proof of Claim 4.2, each component of $F \cap M_{thick}$ is incompressible and meridionally incompressible in $(M_{thick}, T \cap M_{thick})$. Hence, by Lemma 3.3, $F \cap M_{thick}$ has no critical point or F is ∂ -parallel in $(M_{thick}, T \cap M_{thick})$.

In the latter case, if $F \cap M_{thick}$ is ∂ -parallel into S , then F can be isotoped so that it is entirely contained in M_{thin} . Since $(M_{thin}, T \cap M_{thin})$ is a trivial tangle, F is compressible or meridionally compressible in $(M_{thin}, T \cap M_{thin})$. This contradicts the incompressibility or meridionally incompressibility of F in $(B, T \cap B)$. Otherwise, $F \cap M_{thick}$ is ∂ -parallel into ∂B , and we obtain the conclusion 1 of Lemma 3.4.

In the former case, $F \cap M_{thick}$ has no critical point and for any level 2-sphere P in M_{thick} , any loop of $F \cap P$ separates four points of $T \cap P$ into two points and two points in P . It follows that $F \cap M_{thick}$ is a vertical annulus disjoint from T . Hence, F is a disk separating two strings of T , and we have the conclusion 2 of Lemma 3.4. $\square$

4. PROOF

Proof. (of Theorem 2.1) First, we focus the thin regions $M_{thin} = M_1 \cup M_2 \cup \dots \cup M_{k-1} \cup M_k$, where M_1 is the top 3-ball, $M_2, \dots, M_{k-1}$ are thin regions which contain thin level spheres $S_2, \dots, S_{k-1}$ respectively, and M_k is the bottom 3-ball, where $h(S_i) > h(S_{i+1})$ for $i = 2, \dots, k-2$. Put $S_1 = h^{-1}(1 - \epsilon)$ and $S_k = h^{-1}(-1 + \epsilon)$ for a sufficiently small positive real number ϵ . Then, for each i , S_i separates M_i into two submanifolds M_i^+ and M_i^- , where M_i^+ is the upper part and M_i^- is the lower part. Put $K \cap M_i^\pm = \alpha_i^\pm \cup \beta_i^\pm$, where $\alpha_i^\pm$ is a union of arcs each of which has only one critical point, and $\beta_i^\pm$ is a union of vertical arcs. See Figure 14.




FIGURE 14. $M_i^\pm$, S_i , $\alpha_i^\pm$, $\beta_i^\pm$

Claim 4.1. By an isotopy of F in (S^3, K) , we may assume that each component of $F \cap M_i^\pm$ is either an incompressible disk in $(M_i^\pm, K \cap M_i^\pm)$ with only one critical point or an incompressible vertical annulus in $(M_i^\pm, K \cap M_i^\pm)$ disjoint from K . Moreover, if F is meridionally incompressible in (S^3, K) , then we may assume that each component of $F \cap M_i^\pm$ is meridionally incompressible in $(M_i^\pm, K \cap M_i^\pm)$.

Proof. (of Claim 4.1) By an incompressibility of F in (S^3, K) , we may assume that $F \cap S_i$ consists of essential loops in $S_i - K$. Let $T_i^\pm \subset M_i^\pm$ be a union of mutually disjoint monotone arcs connecting a minimal/maximal critical point of $K \cap M_i^\pm$ to a point in $S_i - (F \cap S_i)$. We may assume that F and $T_i^\pm$ are in general position. Put $N_i^\pm = N(S_i \cup T_i^\pm; M_i^\pm)$. Then, each component of $F \cap N_i^\pm$ is either an incompressible disk in $(N_i^\pm, K \cap N_i^\pm)$ or an incompressible annulus in $(N_i^\pm, K \cap N_i^\pm)$ disjoint from K . Since $(M_i^\pm - \text{int} N_i^\pm, K \cap (M_i^\pm - \text{int} N_i^\pm))$ admits a product structure, there exists an isotopy of $\partial N_i^\pm - S_i$ into $\partial M_i^\pm - S_i$ in $(M_i^\pm, K \cap M_i^\pm)$ keeping such conditions. Hence, we have that each component of $F \cap M_i^\pm$ is either an incompressible disk in $(M_i^\pm, K \cap M_i^\pm)$ with only one critical point or an incompressible vertical annulus in $(M_i^\pm, K \cap M_i^\pm)$ disjoint from K . See Figure 15.

Moreover, suppose that F is meridionally incompressible in (S^3, K) . If there exists a component of $F \cap M_i^\pm$ which is meridionally compressible in $(M_i^\pm, K \cap M_i^\pm)$, then it is a vertical annulus. We choose an innermost vertical annulus A and a meridionally compressing disk D for A in $(M_i^\pm, K \cap M_i^\pm)$. Then, there exists a disk D' in F such that $\partial D' = \partial D$ and $D \cup D'$ bounds a trivial (3-ball, arc)-pair in (S^3, K) . An isotopy of F bringing D' to D in (S^3, K) turns the vertical annulus A to a meridionally compressing disk A' in $(M_i^\pm, K \cap M_i^\pm)$, and it can be pushed out from $M_i^\pm$. Here, if A' enters $M_i^\mp$, then it can be pushed out from M_i entirely. Hence, each component of $F \cap M_i^\pm$ is meridionally incompressible in $(M_i^\pm, K \cap M_i^\pm)$. $\square$




FIGURE 15. $M_i^\pm$, S_i , $T_i^\pm$ and $N_i^\pm$

Hence, $F \cap M_i$ consists of incompressible disks with only one minimal/maximal critical point or vertical incompressible annuli in $(M_i, K \cap M_i)$. Moreover, if F is meridionally incompressible in (S^3, K) , then $F \cap M_i$ is also meridionally incompressible in $(M_i, K \cap M_i)$. Hereafter, we suppose that $|F \cap M_{thin}|$ is minimal up to isotopy of F in (S^3, K) under such conditions, and moreover that the number of critical points of $F \cap M_{thick}$ is minimal among all Morse positions of F . We note

that $|F \cap M_{thin}| \neq 0$ since if $F \subset M_{thick}$, then by Lemma 3.2, F is parallel to a thick level sphere and it is compressible.

Claim 4.2. Each component of $F \cap M_{thick}$ is incompressible and meridionally incompressible in $(M_{thick}, K \cap M_{thick})$.

Proof. (of Claim 4.2) Suppose that there exists a component of $F \cap M_{thick}$ which is compressible in $(M_{thick}, K \cap M_{thick})$. Then, there exists a compressing disk D for $F \cap M_{thick}$ in $(M_{thick}, K \cap M_{thick})$. Since F is incompressible in (S^3, K) , there exists a disk D' in F such that $\partial D' = \partial D$. Then, an isotopy of F from D' to D decreases $|F \cap M_{thin}|$ since D' contains at least one component of $F \cap M_{thin}$. This is a contradiction to the minimality of $|F \cap M_{thin}|$. Finally, in the case that F is meridionally incompressible in (S^3, K) , the proof is similar to the case that F is incompressible. $\square$

By the above, the conditions (1), (2) and (4) in Section 1 are satisfied. Moreover, if $F \cap M_{thick}$ does not contain a maximal or minimal critical point, then the condition (3) is satisfied. Hence F is in an essential Morse position related to K , and we have the conclusion 2 of Theorem 2.1.

Hereafter, we suppose that $F \cap M_{thick}$ has a maximal or minimal critical point, and we will show that F is isotopic to a thin level sphere, which is the conclusion 1 of Theorem 2.1. Without loss of generality, let a_m be the lowest maximal critical point of $F \cap M_{thick}$ in a thick region. Then, by Lemma 3.2, a component F_{max} of $F \cap M_{thick}$ containing a_m is ∂ -parallel in $(M_{thick}, K \cap M_{thick})$. If F_{max} is parallel into ∂M_k , then F is isotoped so that it is entirely contained in the trivial tangle $(M_k, K \cap M_k)$, and it is compressible in the trivial tangle. Therefore, without loss of generality, we may assume that F_{max} is isotopic to a planar surface contained in $\partial M_i^+ - S_i$ for a thin region M_i . Put $F \cap M_i^\pm = D_i^\pm \cup A_i^\pm$, where $D_i^\pm$ is a union of incompressible disks with only one critical point and $A_i^\pm$ is a union of vertical annuli in $(M_i^\pm, K \cap M_i^\pm)$. Since each arc of $\alpha_i^\pm$ has only one critical point, there exists a union $\Delta_i^\pm$ of vertical disks in $M_i^\pm$ for $\alpha_i^\pm$ such that $\Delta_i^\pm \cap \alpha_i^\pm = \partial \Delta_i^\pm \cap \alpha_i^\pm = \alpha_i^\pm$ and $\gamma_i^\pm = \partial \Delta_i^\pm - \text{int} \alpha_i^\pm \subset \partial M_i^\pm - S_i$. Moreover, by a horizontal isotopy, we can take $\Delta_i^\pm$ so that $\Delta_i^\pm \cap (D_i^\pm \cup A_i^\pm) = \emptyset$ since each component of $D_i^\pm$ has only one critical point and $A_i^\pm$ is vertical. See Figure 16.


Now, we isotope F_{max} so that it is contained in $\partial M_i^+ - S_i$. If F_{max} contains a component of γ_i^+ , then F_{max} is α_i^+ -compressible and hence F is K -compressible. This contradicts the incompressibility of F in (S^3, K) and we have $F_{max} \cap \Delta_i^+ = \emptyset$. Let F'_{max} be a component of $F_{max} \cup D_i^+ \cup A_i^+ \cup A_i^-$ containing F_{max} and push it into M_i slightly. Since $\Delta_i^\pm \cap (F_{max} \cup D_i^\pm \cup A_i^\pm) = \emptyset$, we may assume that F'_{max} is contained in $M'_i = M_i - \text{int} N(\Delta_i^\pm; M_i)$. We note that $(M'_i, K \cap M'_i)$ admits a product structure $(M'_i, \beta_i^+ \cup \beta_i^-)$.

There are following two cases for F'_{max} .

Case 1: $\partial F_{max} \cap A_i^+ = \emptyset$

Case 2: $\partial F_{max} \cap A_i^+ \neq \emptyset$

In Case 1, F is a 2-sphere which is entirely contained in $(M'_i, \beta_i^+ \cup \beta_i^-)$. By Lemma 3.2, F is ∂ -parallel in $(M'_i, \beta_i^+ \cup \beta_i^-)$. It follows that F is isotopic to a thin level sphere S_i , and we have the conclusion 1 of Theorem 2.1.

FIGURE 16. $M_i^\pm$, S_i , $\alpha_i^\pm$, $\beta_i^\pm$, $\gamma_i^\pm$, $D_i^\pm$ and $A_i^\pm$

In Case 2, F'_{max} is a horizontally ∂ -parallel planar surface in $(M'_i, \beta_i^+ \cup \beta_i^-)$. By Lemma 3.1, F'_{max} is ∂ -parallel in $(M'_i, \beta_i^+ \cup \beta_i^-)$. We isotope F'_{max} in $(M'_i, \beta_i^+ \cup \beta_i^-)$ so that it is contained in M_i^- . Then, there exists a union δ of vertical ∂ -compressing disks for F'_{max} in $(M_i^-, K \cap M_i^-)$ such that $\partial\delta$ cuts F'_{max} into a single disk, and $\delta \cap (D_i^- \cup \Delta_i^-) = \emptyset$. By a ∂ -compression of F'_{max} along δ and a suitable isotopy, F'_{max} turns into a disk with only one maximal critical point in M_i^- . During the transformation above, $|F \cap M_i|$ does not increase, but at least one maximal critical point a_m of $F \cap M_{thick}$ is eliminated. This contradicts the supposition of the number of critical points of $F \cap M_{thick}$. Hence, Theorem 2.1 is proved. $\square$

Proof. (of Theorem 2.3) Let (B, T) be a Montesinos tangle $T(r_1, r_2)$ with $r_i \neq \infty$ for $i = 1, 2$, and D a disk dividing (B, T) into two rational tangles (B_1, T_1) and (B_2, T_2) so that $D = B_1 \cap B_2 = \partial B_1 \cap \partial B_2$ and (B_i, T_i) has a slope r_i . Let F be a closed incompressible surface properly embedded in $B - \text{int}N(T)$. If F is meridionally compressible in (B, T) , then we perform meridionally compressions as possible and obtain a disjoint union F' of closed incompressible and meridionally incompressible surfaces. Conversely, F is obtained from F' by meridional tubings along T . Note that $F' \subset \text{int}B$ since F is a closed surface properly embedded in $B - \text{int}N(T)$.

Claim 4.3. Let S be a closed incompressible and meridionally incompressible surface in (B, T) disjoint from T or intersecting T transversely. Then S is ∂ -parallel in (B, T) .

Proof. (of Claim 4.3) Suppose that $|S \cap D|$ is minimal up to isotopy of S . Then, by the incompressibility and meridionally incompressibility of S in (B, T) , each component of $S \cap B_i$ is incompressible and meridionally incompressible in (B_i, T_i) . We note that $S \cap D \neq \emptyset$ since a rational tangle contains no closed incompressible and meridionally incompressible surface, and that any loop of $S \cap D$ is parallel to ∂D in $D - T$. By Lemma 3.4, each component P of $S \cap B_i$ is ∂ -parallel in (B_i, T_i) or a disk separating two strings of T_i . If there exists a component P of $S \cap B_i$ of the latter type, then $r_i = \infty$ and this contradicts the hypothesis for r_i . Hence, by the minimality of $|S \cap D|$, both of $S \cap B_1$ and $S \cap B_2$ consist of ∂ -parallel disks that are not parallel into D . By connecting these components of $S \cap B_i$, we have a 2-sphere S which is ∂ -parallel in (B, T) . $\square$

By Claim 4.3, F' is a ∂ -parallel 2-sphere in (B, T) . Now, we recover F from F' by tubing F' along T . Then F is ∂ -parallel in $B - \text{int}N(T)$, hence the Montesinos tangle $T(r_1, r_2)$ is small. $\square$

Remark 4.4. In general, Montesinos tangles are not small. Tsutsumi informed me that there are Montesinos tangles with length more than two that are not small. Actually, any closed incompressible and meridionally incompressible surface in a Montesinos tangle $T(r_1, \dots, r_n)$ are parallel to $\partial(B_i \cup \dots \cup B_j)$, where $B_i, \dots, B_j$ are successive two or above rational tangles. Then, we perform a meridional tubing along $T_i \cup \dots \cup T_j$ and obtain a closed incompressible surface which is not ∂ -parallel in the exterior of $T(r_1, \dots, r_n)$.

Proof. (of Theorem 2.6) Let G be a 2-bridge theta curve or handcuff graph, F an incompressible and meridionally incompressible closed surface in (S^3, G) . In the same way as the case of knots, we decompose S^3 into $M_{thin} = M_1 \cup M_2$ and M_{thick} , where M_1 is the top 3-ball and M_2 is the bottom 3-ball. Then, one vertex and the maximal critical point of G are contained in M_1 , and another vertex and the minimal critical point of G are contained in M_2 . Moreover, by the proof of Theorem 2.1, we may assume that $F \cap M_1$ consists of incompressible disks in $(M_1, G \cap M_1)$ each of which has only one maximal point and disjoint from G , $F \cap M_{thick}$ consists of incompressible and meridionally incompressible surfaces each of which has no minimal/maximal critical point, $F \cap M_2$ consists of incompressible disks in $(M_2, G \cap M_2)$ each of which has only one minimal point and disjoint from G . Suppose that $|F \cap M_{thin}|$ and the number of critical points of $F \cap M_{thick}$ is minimal up to isotopy of F in (S^3, G) . By Lemma 3.3, $F \cap M_{thick}$ has no critical point or F is ∂ -parallel in $(M_{thick}, G \cap M_{thick})$.

In the latter case, F can be isotoped so that it is entirely contained in M_{thin} . Then, F is compressible or meridionally compressible in $(M_{thin}, G \cap M_{thin})$ since $G \cap M_{thin}$ is contained in a disk properly embedded in M_{thin} . This is a contradiction.

In the former case, $F \cap M_{thick}$ has no critical point and for any thick level 2-sphere P for G , any loop of $F \cap P$ separates five points of $G \cap P$ into two points and three points in P . It follows that $F \cap M_{thick}$ is a vertical annulus, and F is a sphere with one maximum and one minimum. Hence, F is a 2-sphere illustrated in Figure 6. $\square$

Proof. (of Theorem 2.8) Let L be a link in S^3 which admits a Hopf tangle decomposition $(S^3, L) = (B_1, T_1) \cup (B_2, T_2)$, and F be an incompressible and meridionally incompressible surface in (S^3, L) . For each $i = 1, 2$, the trivial loop $C_i \subset T_i$ bounds a disk D_i in B_i which intersects each string of T_i in only one point. Suppose that $|F \cap (D_1 \cup D_2)|$ is minimal up to isotopy of F . By the incompressibility and meridional incompressibility of F , each component of $F \cap D_i$ is either an arc separating two points of $T_i \cap D_i$ in D_i or a loop parallel to ∂D_i in $D_i - T_i$. Put $H_i = N(D_i)$. Then, $(H_i, T_i \cap H_i)$ is also a Hopf tangle and each component of $F \cap H_i$ is either a “meridian” disk bounded by the meridian of $(H_i, T_i \cap H_i)$ or a “vertical” annulus whose boundary component is parallel to the equator of $(H_i, T_i \cap H_i)$ in $\partial H_i - \partial(T_i \cap H_i)$. See Figure 17. We remark about $F \cap H_i$ that a meridian disk is not compatible with a vertical annulus.




FIGURE 17. a “meridian” disk and a “vertical” annulus

The remainder of these two small Hopf tangles $S^3 - \text{int}(H_1 \cup H_2)$ admits a product structure $(S \times I, X \times I)$, where S is the decomposing sphere for (S^3, L) and X is four points of $L \cap S$. Under the minimal condition of $|F \cap (D_1 \cup D_2)|$, we put $F \cap (S \times I)$ in a Morse position with respect to the height function $h : S \times I \rightarrow I$ and suppose that the number of critical points of $F \cap (S \times I)$ is minimal among all Morse positions of $F \cap (S \times I)$. In the same way as Claim 4.2, each component of $F \cap (S \times I)$ is incompressible and meridionally incompressible in $(S \times I, X \times I)$. It follows from Lemma 3.3 that each component of $F \cap (S \times I)$ is either a vertical annulus in $(S \times I, X \times I)$ which separates four strings of $X \times I$ into two and two, or ∂ -parallel in $(S \times I, X \times I)$. In the latter case, F is parallel to the decomposing sphere S when $F \cap (D_1 \cup D_2) = \emptyset$, or it contradicts the minimality of $|F \cap (D_1 \cup D_2)|$ or the incompressibility and meridional incompressibility of $F \cap (S \times I)$. In the former case, L belongs to $\mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2) \cup \mathcal{L}(e_1, e_2)$ since any loop of $F \cap \partial H_i$ is either the meridian or the equator of $(H_i, T_i \cap H_i)$ and each component of $F \cap (S \times I)$ is a vertical annulus disjoint from $X \times I$. Thus the conclusion 1 of Theorem 2.8 holds.

Next, if $L \in \mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2)$, then there exists another Hopf tangle decomposing sphere which is made of meridian disks and vertical annuli in H_i and vertical annuli in $S \times I$ as Figure 9. Conversely, if there exists a Hopf tangle decomposing sphere different from S , then it is an incompressible and meridionally incompressible closed surface in (S^3, L) . Hence, by the conclusion 1 of Theorem 2.8, we have $L \in \mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2) \cup \mathcal{L}(e_1, e_2)$. But for a link

which belongs to $\mathcal{L}(e_1, e_2) - \mathcal{L}(m_1, m_2)$, there does not exist an another Hopf tangle decomposing sphere, therefore we have $L \in \mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2)$. Thus the first sentence of conclusion 2 of Theorem 2.8 holds. Moreover, since any Hopf tangle decomposing sphere different from S is made of meridian disks and vertical annuli in H_i and vertical annuli in $S \times I$ as Figure 9, $L \in \mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2)$ has exactly two Hopf tangle decompositions, except the Borromean rings has exactly three Hopf tangle decompositions as Example 2.9. Thus the conclusion 2 of Theorem 2.8 holds.

Finally, suppose that L is not small and let Q be an incompressible and not ∂ -parallel closed surface in $E(L)$. Then, by meridional compressing Q as possible, we get an incompressible and meridionally incompressible closed surface Q' in (S^3, L) . By the conclusion 1 of Theorem 2.8, we have $L \in \mathcal{L}(m_1, m_2) \cup \mathcal{L}(m_1, e_2) \cup \mathcal{L}(e_1, m_2) \cup \mathcal{L}(e_1, e_2)$. But if $L \notin \mathcal{L}(e_1, e_2)$, then any meridional tubing Q' cannot be performed since Q' must be a Hopf tangle decomposing sphere. Hence, $L \in \mathcal{L}(e_1, e_2)$ and Q is an essential torus illustrated in Figure 9. Thus the conclusion 3 of Theorem 2.8 holds. $\square$

The following exercise can be proved by the similar method as Lemma 3.4. So, we omit the proof.

Exercise 4.5. Let (B, T) be a Hopf tangle and F a surface properly embedded in B which is disjoint from T or intersects T transversely. If F is incompressible and meridionally incompressible in (B, T) , then one of the following holds.

- (1) F is a meridian disk for (B, T) .
- (2) F is a vertical annulus for (B, T) .
- (3) F is ∂ -parallel in (B, T) .

Acknowledgement. The author would like to thank Yukihiro Tsutsumi for informing Remark 4.4, and the referee for pointing out the third Hopf tangle decomposing sphere for the Borromean rings.

REFERENCES

1. D. Gabai, Foliations and the topology of 3-manifolds III, J. Diff. Geom. **26** (1987), 479–536.
2. C. McA. Gordon and J. Luecke, Knots are determined by their complements, J. Amer. Math. Soc. **2** (1989), 371–415.
3. M. P. Heegaard, *Sur l'Analysis Situs*, Bull. Soc. Math. France **44** (1916), 161–242.
4. H. Matsuda, *Small knots and links*, preprint.
5. T. Motohashi, *2-bridge θ -curves in S^3* , Topology Appl. **108** (2000), 267–276.
6. U. Oertel, *Closed incompressible surfaces in complements of star links*, Pacific J. Math. **111** (1984), 209–230.
7. H. Schubert, *Über eine numerische Knoteninvariante* Math. Z. **61** (1954), 245–288.
8. Y.-Q. Wu, *The classification of nonsimple algebraic tangles*, Math. Ann. **304** (1996), 457–480.

DEPARTMENT OF NATURAL SCIENCES, FACULTY OF ARTS AND SCIENCES, KOMAZAWA UNIVERSITY, 1-23-1 KOMAZAWA, SETAGAYA-KU, TOKYO, 154-8525, JAPAN

E-mail address: w3c@komazawa-u.ac.jp